

In-vivo PIV Measurement of the Blood Flow in the Heart and Main Vein of Medaka Embryo

Nao Ninomiya^{1*}, Masaru Matsuda², Natsuo Araki¹, Narumi Yano¹

1: Dept. of Opt. Eng., Utsunomiya University, Japan

2: Center for Bioscience Res. and Edu., Utsunomiya University, Japan

* Correspondent author: nino@utmu.jp

Keywords: In-vivo Measurement, PIV, RBC, Heart, Main Vein, Medaka Embryo

ABSTRACT

In order to examine the generation mechanism of the vertebrates, including humans, and to explore the diverging point of the genetic abnormalities, it is effective to investigate the growth of the embryo of an artificial mutant. As the embryos of Medaka hatch outside of the body and their egg membrane are transparent, it is quite easy to carry out the in-vivo observation of their organs. Moreover, the development and the structure of their major organs of Medaka are quite similar to those of human. Thus, the observation of the developmental abnormalities of Medaka embryos may serve as a clue to the pathologic investigation of the genetic disease and hopefully give some insight to the future effective treatment. Presently, the intracardiac blood flows of Medaka embryos have been observed by a high-speed camera under microscope. The obtained images are pre-processed and the velocity fields in the heart and in the main vein of Medaka embryo are measured by PIV method. The red blood cells flowing inside the heart and in the main vein of Medaka embryos are used as tracer particles. The image pre-processing was applied to capture the image of red blood cells and then the cross-correlation method can be applied. As a result, it is found that the velocity defect caused by the valve closing is clearly observed in the main vein and the amplitude of the velocity defect is decreasing with the distance to the heart. It is also found that the beating of the heart corresponds to the pulsatio of the flow velocity of the red blood cell in the heart. In conclusion, the authors observed the Medaka embryo under microscope with a high-speed camera and measured the velocities of the red blood cells in the heart and in the main vein by PIV measurement using the pre-processed images.

1. Introduction

In order to examine the generation mechanism of the vertebrates, including humans, and to explore the diverging point of the genetic abnormalities, it is effective to investigate the growth of the embryo of an artificial mutant. As the embryos of Medaka or Zebrafish hatch outside of the body and their egg membrane are transparent, it is quite easy to carry out the in-vivo observation of their organs. Moreover, the development and the structure of their major organs of Medaka are quite similar to those of human (Omran et al. 2008). The DNA of Medaka has been already read and it is easy to cause the genetic abnormalities to the mutant. Thus, the observation of the developmental abnormalities of Medaka embryos may serve as a clue to the pathologic

investigation of the genetic disease and hopefully give some insight into the future effective treatment.

2. Visualization

Presently, the in-vivo intracardiac blood flows of Medaka embryos have been observed by a high-speed camera under microscope as shown in Fig. 1. The microscope used in this study is Leica Microsystems MZ16 and the high-speed camera is Photron FASTCAM Mini WX100, working frame rate is 125fps with the shutter speed of 1/1000 sec is selected and the resolution is 2048 x 2048.




Fig. 1 Experimental setup.

The obtained images are processed and the velocity vectors in the heart and in the main vein are measured by PIV. In this study, red blood cells flowing inside of the heart of Medaka embryos are used as tracer particles, but the images of red blood cells look dark in the captured image as shown in Fig. 2(a), so PIV measurement cannot be performed with the images of the red blood cell as it is. Therefore, image pre-processing has been applied to the images. First, the brightness of the image is inverted as shown in Fig. 2(b) so that the red blood cell can be captured as the bright fragments as is the case for the tracer particles. Second, the brightness of the darkest instance of each pixels has been collected to obtain the background brightness as shown in Fig. 2(c). Finally, by subtracting the background brightness, only the image of the red blood cell is highlighted and thus the cross-correlation method commonly used by PIV can be applied.




Fig. 2 Pre-processing to the image.

3. PIV Measurement

The typical result of the PIV in the main vein is shown in Fig. 3. The color of the velocity vectors indicates the velocity magnitude from blue to red and the velocities magnitudes are shown by $\mu\text{m/s}$.




Fig. 3 PIV results in the main vein.

The typical result of the PIV in the heart is shown in Figs. 4. The PIV results are super-imposed onto the original image of the Medaka embryo together with the manually drawn outline of the heart. The outline of the atrium is drawn by red, the valve by yellow and the ventricle by green.




Fig. 4 PIV results in the heart.

As the embryo of Medaka is spherical and the organs inside of the embryo is located three-dimensionally, the embryo has been placed upon an agar culture medium so that a good focus can be obtained on the main vein or the heart as shown in Figs. 2. The size of the interrogation window is set as 32×32 and the search regions are set as 16×16 for the main vein and 32×32 for the heart, and the overlaps are 50% and 75% respectively. It can be seen that the velocity vectors in the main vein or in the heart are clearly obtained by cross-correlation algorithm of PIV with the aid of pre-processing to the images obtained by the microscopy and the high-speed camera.

The typical size of the embryo is 1 mm and the typical width of the main vein is about $50 \mu\text{m}$ and the typical velocity in the main vein is about $700 \mu\text{m/s}$. As the blood in the Medaka embryo is dilute, it may be feasible to assume the physical properties of the blood is almost same as those of normal saline solution. Thus, the Reynolds number in the main vein is estimated as 0.035 and thus the flow is laminar.




Fig. 5 Time variation of the velocities in the main vein.

Figure 5 shows the time variation of the velocities in the main vein at different locations close to the inlet of the heart from A to E, where E is closest to the heart. The velocities at the center of the main vein are phase-averaged by synchronizing the velocity changes by the velocity minimum in each heart beat and the 20 cycles are averaged to obtain the time variation during an averaged cycle of 0.48 sec.

Due to the continuity constraints, the velocity decreases with the increase in the diameter of the main vein. Moreover, negative pulses in the velocity are observed according to the heart beat. One interesting finding is that the pulse in the velocity does not have a time delay along the main vein. Assuming that the blood is incompressible, it suggests that the change in the diameter of the main vein is very small. Another interesting finding is that the magnitude of the velocity defects due to the heart beat decreases with the distance to the heart. This implies that the membrane of the main vein has a compliance to reduce the velocity change.

Figures 6 show the PIV results in the atria and in the end point of the main veins. Three main veins merge into one at the inlet of atria. The outline of the atria is drawn by red and those of the main veins by pale blue.




Fig. 6 PIV results in the atria of the heart.




Fig. 7 Time variation of the velocities in the atria.

The time variation of the velocity in the atria is also obtained as shown in Fig. 7. The phase average is obtained by synchronizing the sharp velocity decrease at the inlet of the atria, which is shown by purple dot in Fig. 6 and labeled by [A] in Fig. 7. One cycle of the embryo at this age was 0.60 sec. The typical age of the embryo used in this study is about 2 days after birth.

It is found that there is no valve at the inlet of the atria of the Medaka heart and a pyloric structure at the inlet of the heart controls the flow direction. The blood flows as if a man shuts the mouth and sucks and swallows the liquid into the palate. As a result, the movie of the shape changes of the organs together with the velocity vectors of the red blood cell is prepared. According to the movie, it is found that the beating of the heart corresponds to the pulsation of the flow velocity of the red blood cell in the heart.

The measurement of the velocity in the ventricle has also been tried, but it was very difficult to see the red blood cells in the ventricle because the ventricle is located close to the eyeballs.

Consequently, the authors observed the blood flow in the heart of Medaka embryo under microscope with a high-speed camera and measured the in-vivo velocities of the red blood cells in the heart by PIV measurement using the pre-processed images.

4. Future Study

Presently, the in-vivo velocity measurement in the heart and in the main vein of Medaka embryo has been carried out and the velocity change in the heart has been investigated along with the shape change of the atrium. In the future study, by comparing the velocities in the healthy embryo and those in the embryo with the heart disease that has been artificially caused by DNA manipulation, the early detection of the developmental abnormalities may be possible and it may serve as a clue to the pathologic investigation of the genetic disease and hopefully give some insight to the future effective treatment at early stage.

The present PIV measurement has been carried out in the two-dimensional plane which is parallel to the blood flow in the heart of the Medaka embryo and thus the velocity information obtained in this study has been limited to two-dimensional information. But, in order to fully investigate the flow conditions in the heart, three-dimensional velocity measurement should be carried out. The authors of this study have invented the three-dimensional velocity measurement by using Doppler phase-shifting holography which can be used with single camera observation (Ninomiya et al. 2016). The three-dimensional velocity measurement in the heart of the Medaka embryo using Doppler phase-shifting holography remains for the future study.

5. Conclusions

The in-vivo velocity measurement of the blood flow in the heart of Medaka embryo has been carried out. The images of the red blood cell in the heart of the Medaka embryo have been taken by the high-speed camera and the microscope. The images have been pre-processed to highlight only the image of the red blood cell and to enable the PIV measurement. Consequently, the phase-averaged velocity changes in the vena cava and in the atria has been obtained. The results help in finding the developmental abnormalities in the early stage and may give some insight to the future effective treatment.

6. Permissions

The authors hold all the rights to publish everything in this manuscript.

Acknowledgements

A part of this work has been supported by UU Interdisciplinary Project for in-vivo Bioimaging and Sensing from MEXT and the JSPS KAKENHI Grant Number 16K06107. All supports are greatly acknowledged.

References

- Omran, H., Kobayashi, D., Olbrich, H., Tsukahara, T., Loges, N. T., Hagiwara, H., Zhang, Q., Leblond, G., O'Toole, E., Hara, C., Mizuno, H., Kawano, H., Fliegau, M., Yagi, T., Koshida, S., Miyawaki, A., Zentgraf, H., Seithe, H., Reinhardt, R., Watanabe, Y., Kamiya, R., Mitchell, D. R., and Takeda, H., Ktu/PF13 is required for cytoplasmic pre-assembly of axonemal dyneins, *Nature* 256, pp. 611-616 (2008)
- Ninomiya, N., Kubo, Y., Barada, D. and Kiire, T., Three-Dimensional Velocity Measurement by a Single Camera using Doppler Phase-Shifting Holography, *Measurement, Science and Technology*, Vol. 27, No. 10, 104004 (2016)